1. C. gynocrates, Wormsk. (C. disica 1st Edn now L.)
2. Boott would prefer this removed to the "Stellulatæ", to follow C. stellulata, & it may perhaps belong there, of right, but this Division "Psyllophoræ," as here constituted, professes to be merely "artificial"—
6½ xxx. leaves very broad = longer than the culm C. Fraseri.
11. (C. Liddoni, 1st Edn now Boott!)
12 scarcely to be distinguished from C. intermedia, Good.
14 Boott thinks that this species should rather be referred as a var. of the last (C. teret.) but I think the perig.ne is clearly that of C. paniculata = indeed I am convinced of it.

19. Kentucky. Short! in Herb.
Boott thinks this is probably only a var. of C. stipata, but this I doubt.

20. Washington, Mich.n Dr. Cooley & Bois Blanc Island. Detroit River - Dr. P. H. Maclagan.

21. Lexington Kent.y. Short! (Cincinnati Lea! both in Herb. Torrey. — Closely spiked heads seem too near No. 20, tho' the perig.ia appear always larger & more tapering.

28. C. tenella. Schk.r now Fbrh.
(C. gracilis 1st Edn now Fbrh. fide Boott! ex spec. orig.!) N.B. Boott has ascertained that C. gracilis of Fbrh. is the true C. foliacea Lin.

from an authentic spec.ⁿ in Herb. Smith. —
C. tenella Fbrh. is C. canescens, L.

31. Var. 2. is C. vitilis, Fries. ex Boott! & the Syn. may be quoted. (see Boott's letter)

35. Detroit river — Dr. Maclagan!

36. Omit this species, not in the U. S. (see Syn. No. 11)

38½. C. adusta — Boott. Rhode Isld — Olney — Catskill Mt. L. C. Lake Mahopac. NJ. J. C. — & Lake Superior. Loring!

43. This is C. verrucosa, Schweinitz! in Herb. Torrey! Mountain runs — Clinton Co. Penn.ª J. M. McMinn Mts of Nor. Car. Buckley.

46. Lamark's name must stand for this species, as he founded it on an Amer.ⁿ plant. & Goodenough's C. stricta has never been found in America.

47½. C. lenticularis, Mx. Lake Avalanche. Torr. in Herb. Boott! N. Shore of Lake Superior. Loring Jr.!

48½. C. maritima Mütt. In Herb. Torrey are forms just resembling the foreign from N. Hamp.ª J. Lesquereux Massachusetts. Oakes, & N. J. P. D. Sr.!

49. Var. 1. C. crinita, muricata. J. C. in Silt. Journal. (C. paleacea 1st Edn non Wahl.)

54½. C. Vahlii, Schk. Isle Royal Whitney! Pic River. C. Loring Jr.!

59. A specimen of what I take for C. Wormskiöldii is in Herb. Torrey! from Macon Co. N. C. Buckley. Dec.

61. After Syn. C. heterostachya Torr. non Dew.

68½. C. obstipata. Curtis. "Side of Saddle Mount. Haf. 800. ft. above the base." Dewey. Herb. Torr.! also Pokono Mt. Pa. Darlington & Toorus. — & West Chester. Pa. Townsend.

72. Border of a marsh — near Bristol. Penn.ª Dr. Eberle in Herb. Short! after authority — C. Careyana Torr non Dew.

76. C. lapiflora, Lam. non Schk. (C. laceps with. & of 1st Edn.)

85½. C. Richardsonii Hk. near Rochester, N. Y. "within a few miles of the city" Dew. to A. Gray 15 Feby 1848. Prairies Imputa, St. Mead — Macomb Co. Michigan. Coley.

90. Montreal. Dr. Maclagan.

93½. C. Cherokeensis, Schwz. & Torr. — near Natchez. Miss. W. S. Sullivant!

94. The Syn. C. Greeniana belongs here & that of C. binervis to 95 very spec.ⁿ in Herb. Torr.! —

101. Boott considers that in justice to Muhl. his name of C. polymorpha ought to attach

to this species (to which it belongs in part)
restricting C. striata Mx. for No. 102.

102. C. striata, Mx. (not of 1st Edn.)

104. Spoken to Saddle Mountain Kooskooskie river -
U. S. Explg. Expdn!

105. C. trichocarpa. Muhl. (C. striata, 1st Edn. not of Mx.)

107½ C. media, Dew. Greece. Monroe Co. N. Y. Dr. Bradley -
a single spec.n in Herb. Torr! & Sartwell has
another - see them for the description, as the
species is very remarkable & only found, I
believe, by Dr. Bradley, who doubtless has at
least one specimen himself, as Sartwell derives
his from him - Dewey (in Woods Bot.y) confounds
2 species, that from Georgia being totally
distinct - In his Carices I he formerly
referred Bradley's plant to C. gigantea -
vide Herb. Torr!

114. South Kingston - Rhode Island. Olney!

115. The var. polystachya is Sartwell's C. lupuliformis -
if you think the characters he gives sufficient.

118½ C. gigantea, Rudge - Banks of the Ohio, at North Bend -
& near Louisville. Short! also Florida Chapman!
& Texas Wright &c. - well distinguished from C. lupulina
by the less tumid, horizontally spreading perig.a & many male spikes.